

8-21-2013

Age- and Experience-Related Distribution of Parvalbumin-Expressing GABAergic Interneurons within the Hippocampus & Surrounding Cortices

Jennifer A. Corriveau

University of Connecticut - Storrs, jennifer.corriveau@uconn.edu

Recommended Citation

Corriveau, Jennifer A., "Age- and Experience-Related Distribution of Parvalbumin-Expressing GABAergic Interneurons within the Hippocampus & Surrounding Cortices" (2013). *Master's Theses*. 489.
https://opencommons.uconn.edu/gs_theses/489

This work is brought to you for free and open access by the University of Connecticut Graduate School at OpenCommons@UConn. It has been accepted for inclusion in Master's Theses by an authorized administrator of OpenCommons@UConn. For more information, please contact opencommons@uconn.edu.

Age- and Experience-Related Distribution of Parvalbumin-
Expressing GABAergic Interneurons within the Hippocampus &
Surrounding Cortices

Jennifer Ann Corriveau

B.A., Colby College, 2010

Master of Arts Thesis

Submitted in Partial Fulfillment of the

Requirements for the Degree of

Master of Arts

At the

University of Connecticut

2013

APPROVAL PAGE

Master of Arts Thesis

Age- and Experience-Related Distribution of Parvalbumin- Expressing GABAergic Interneurons within the Hippocampus & Surrounding Cortices

Presented by:

Jennifer Ann Corriveau, B.A.

Major Advisor: _____

Dr. James J. Chrobak, Ph.D.

Associate Advisor: _____

Dr. John D. Salamone, Ph.D.

Associate Advisor: _____

Dr. Chi-Ming Chen, Ph.D.

University of Connecticut

2013

Acknowledgements

I would like to extend my sincerest gratitude to my graduate advisor and primary investigator, Dr. James J. Chrobak, for his continued support, advice, knowledge, and patience during the entirety of my time at UConn thus far, but particularly throughout the research and preparation of my Master of Arts thesis. I would also like to thank my committee members, Dr. John D. Salamone and Dr. Chi-Ming Chen, for all of their advice, conversations, and consideration during this process. In addition, I could not have completed this work without the support of my lab-mate, Lauren Long, M.A., and my fellow BNS graduate students: Nicholas Bannon, M.A.; Dr. Eric Nunes; and Patrick Randall, M.A., who have graciously provided me with moral and technical support on this project. Last, but certainly not least, I would like to thank my parents, Ron and Ann Corriveau, as well as my fiancée, Christina Honeycutt, for their continued love and support in all that I do.

Table of Contents

| | |
|---|----|
| Chapter One: Introduction | 1 |
| Goals of Current Research | 1 |
| Schizophrenia | 2 |
| GABAergic Inhibitory Interneurons | 6 |
| Treatment and Age-Related Changes in Parvalbumin Expression | 8 |
| Chapter Two: Manuscript | 11 |
| Abstract | 11 |
| Introduction | 13 |
| Materials and Methods | 15 |
| Results | 19 |
| Discussion | 23 |
| Chapter Three: Discussion | 30 |
| Contributions of the current work to the literature: Age-related changes in PV expression | 30 |
| Contributions of the current work to the literature: Experience-related changes in PV expression | 33 |
| Future Directions | 34 |
| General Conclusions | 37 |
| References | 38 |
| Figures | 50 |
| Tables | 69 |

List of Figures & Tables

Figures

| | |
|--|----|
| Figure 1: <i>Location of representative sections relative to Bregma.</i> | 50 |
| Figure 2: <i>Optimizing PV Antibody Concentration for Visualization of PV-containing Interneurons.</i> | 51 |
| Figure 3: <i>Anatomical locations of photomicrographs taken for analysis of PV expression in hippocampal regions of interest.</i> | 52 |
| Figure 4: <i>Example of PV-expressing interneuron quantification using ImageJ color-based thresholding.</i> | 53 |
| Figure 5: <i>Septotemporal differences in PV expression as a function of hippocampal sub-region and age.</i> | 54 |
| Figure 6: <i>Distribution of HPC CA1 PV+ interneurons as a function of age and septotemporal location.</i> | 55 |
| Figure 7: <i>Distribution of HPC CA3 PV+ interneurons as a function of age and septotemporal location.</i> | 56 |
| Figure 8: <i>Distribution of HPC DG PV+ interneurons as a function of age and septotemporal location.</i> | 57 |
| Figure 9: <i>Age-dependent decrease in parvalbumin expression within the hippocampal sub-regions CA1 and DG, but not CA3.</i> | 58 |
| Figure 10: <i>24-month-old rats have significantly higher PV+ cells within retrosplenial and somatosensory cortices than 1- and 6-month-old rats.</i> | 59 |
| Figure 11: <i>PV+ expression within the retrosplenial cortex across age groups.</i> | 61 |
| Figure 12: <i>PV+ expression within the somatosensory (barrel field) cortex: distribution depends on age.</i> | 62 |

| | |
|---|----|
| Figure 13: <i>Hippocampal CA1 has significantly more PV expression than CA3 and DG in septal and midseptotemporal regions across all age groups.</i> | 64 |
| Figure 14: <i>24-month old extensively trained rats have elevated PV expression across all hippocampal regions.</i> | 65 |
| Figure 15: <i>Proposed age-dependent decline in PV expression as a function of brain region.</i> | 67 |

Tables

| | |
|--|----|
| Table 1. <i>Average number of parvalbumin-positive interneurons across cortical and hippocampal regions in the rat brain as a function of age and septotemporal location.</i> | 69 |
|--|----|

Chapter One: Introduction

Goals of Current Research

The present study sought to elucidate upon the paucity of findings currently available with regard to age-related differences in parvalbumin (PV) expression across brain regions and the septotemporal axis of the hippocampus (HPC) in rats. The lack of available information regarding normative distribution of this calcium binding protein is surprising, given the multitude of studies demonstrating: 1) decreased distribution of PV-expressing GABAergic neurons in post-mortem tissue from schizophrenic patients, and 2) a decrease in PV-expressing GABAergic neurons in forebrain tissue subsequent to acute and chronic NMDA antagonist treatment (e.g., Kittelberger et al., 2012; Wang et al., 2008; Abekawa et al., 2007; Braun et al., 2007; Keilhoff et al., 2004). The latter of which is a widely used model to examine the neuropathological events, as well as the altered behavior/cognitive processes, seen in schizophrenia.

In chapter one, we provide context for the current study, first with a description of the features of schizophrenia and the anatomical changes observed in post-mortem human brain tissue. Secondly, we discuss the importance of observed changes in local circuit GABAergic neurons, and define the anatomical and morphological features of key GABAergic cell populations. Lastly, we discuss the literature regarding treatment- and age-related changes in PV expressing neuron populations.

Schizophrenia

Schizophrenia is a complex psychological disorder affecting approximately 1% of the population worldwide (Perälä et al., 2007). The illness is characterized by positive symptoms including hallucinations, delusions, and paranoia; negative symptoms which include social deficits and flattened affect; and, of particular note, cognitive symptoms which can manifest as marked deficits in attention and memory (Lewis & Lieberman, 2000; Gold, 2004). Currently, positive and, to a certain extent, negative symptoms can be managed through pharmacological treatment with antipsychotic medication such as haloperidol, risperidone, and clozapine, to name a few (Carman et al., 1995; Leucht et al., 2009). However, there are no antipsychotic pharmaceutical interventions capable of mediating the debilitating cognitive deficits characteristic of schizophrenia (Buchanan et al., 2007). Within the past two decades, there has been a concerted effort to understand the underlying etiology of cognitive deficits in schizophrenia. In order to better understand these deficits, and thus foster development of novel cognitive-targeting interventions, it would be helpful to establish a more comprehensive understanding of the neurobiology of forebrain circuits that support the cognitive processes which are altered in schizophrenia.

There are many theories regarding the etiology of schizophrenia, including, but not limited to: genetic predisposition (Bergen & Petryshen, 2012; Kukshal et al., 2012; Owen, 2012; Greenwood et al., 2013), prenatal infection (Brown & Derkits, 2009; Brown & Patterson, 2011), environmental influence (van Os et al., 2010; Bourque et al., 2012), as well as a large body of evidence

supporting combinations of the aforementioned (e.g. Clarke et al., 2009; Holtzman et al., 2012; Pelayo-Terván et al., 2012). However, it is clear from imaging and postmortem studies that distinct anatomical changes – including: volumetric (Bogerts et al., 1985; Honea et al., 2005; Haijma et al., 2012), circuitry (Lewis & Sweet, 2009; Uhlhaas & Singer, 2010), connectivity (Camchong et al., 2011; Schmitt et al., 2011), and neuronal (Harrison, 1999) - contribute to the core cognitive dysfunction in schizophrenia. Importantly, forebrain circuitry, including, but not limited to, prefrontal cortex and temporal lobe structures, are found to have altered regional activity (e.g. magnetic resonance imaging; Shenton et al., 2001; Schobel et al., 2009), differences in white matter connections (e.g. diffusion tensor imaging; Kubicki et al., 2007), and regionally specific volumetric decreases (Sigmundsson et al., 2001) in schizophrenic patients as compared to healthy controls.

Circuitry Implicated in Schizophrenic Cognitive Dysfunction

Cognitive dysfunction is pervasive throughout the lifetime of schizophrenic patients, and severity of cognitive deficits is independent of other psychotic symptoms, and even predictive of prognosis (Tamminga et al., 1998). Much emphasis has been placed on the decreased ability of patients to complete working memory tasks, particularly those involving a high degree of information manipulation (Barch, 2006). Dorsolateral prefrontal cortical (DLPFC) circuitry is implicated in these working memory deficits, and studies of working memory in schizophrenics have even demonstrated an alterations in DLPFC activation in

patients during working memory tasks (Tan et al., 2007). Additionally, hippocampal circuitry, as well as hippocampal-cingulate cortical circuit systems, is thought to be altered at both macro- (e.g. regional interconnections) and micro- (e.g. local inhibition onto pyramids) levels (Benes, 2000). Within these circuits, we are particularly interested in changes seen in GABAergic local circuit networks, as they play a critical role in regulating cognitive and memory processes (Buzsaki & Chrobak, 1995). Furthermore, evidence suggests that there are evolutionary changes within select GABAergic neurons within the forebrain contribute to developmental alterations in cortical and hippocampal macro- and micro-circuits (Hof et al., 1999; Burns, 2004). These alterations can manifest as cognitive dysfunction in a number of psychiatric disorders (e.g., schizophrenia, autism, Alzheimer's; Raghanti et al., 2010).

Alterations in activation seen during neuroimaging studies in DLPFC are likely due to dysfunctional neuronal activity. Specifically, a decrease in GABAergic neurotransmission, and consequently decreased inhibition of excitatory pyramidal neurons, and conversely, even increased inhibition due to compensatory up-regulation of GABA receptors (Gonzalez-Burgos & Lewis, 2008), have been described (Volk & Lewis, 2002; Lisman et al., 2008). Of particular note, the selective decrease of PV-expression within inhibitory GABAergic interneurons, especially within prefrontal cortex and hippocampus, is thought to play a key role in these observed alterations (Lewis et al., 1999; Hashimoto et al., 2003; Lewis et al., 2004). Disrupted signaling within this circuit is likely the driving force behind these deficits, and therefore inhibitory

interneurons are being considered as targets for possible cognitive-enhancing treatments (Lewis et al., 2005).

Afferent and efferent connections between the DLPFC and hippocampus are also thought to be altered, thereby contributing to deficits seen during cognitive processes that require the manipulation and temporal and/or spatial encoding of information (Meyer-Lindenberg et al., 2005). Furthermore, local inhibitory circuitry within the hippocampus has been found to be altered at the synaptic level (e.g. aberrant functionality and synaptic connections) and contributes to widespread cognitive dysfunction (Harrison, 2004). The GABAergic alterations within the prefrontal cortex and hippocampus are further implicated in post-mortem studies of schizophrenic brains, where immuno-reactive labeling studies reveal a selective reduction of the calcium-binding protein parvalbumin, relative to controls (Beasley & Reynolds, 1997; Zhang & Reynolds, 2002). While there is a selective reduction in PV-expressing GABAergic neurons, there is no concurrent decrease in either calretinin or calbindin populations (Zhang & Reynolds, 2002; Lewis et al., 2012). Why forebrain PV-expressing GABAergic neurons would be selectively disrupted is unclear, although evidence presented in the current study suggest developmental changes in PV expression that perhaps make these neurons vulnerable to dysfunction.

GABAergic Inhibitory Interneurons

GABAergic interneurons are integral for optimal functioning within the brain, and play important roles in regulating neurotransmission and coordinating concerted neuronal ensembles (Buzsaki & Chrobak, 1995). These interneurons can be divided into various subtypes, often by the calcium-binding proteins they express (McBain & Fisahn, 2001). Within the brain, there are many different populations of inhibitory interneurons expressing calcium-binding proteins, some of which are used as cellular markers, including: calbindin, calmodulin, calretinin, and parvalbumin (Freund & Buzsaki, 1996; Markram et al., 2004). These calcium-binding proteins allow for the binding and intracellular transportation of calcium ions and can differentially affect the temporal dynamics of cellular function post stimulation (Arif, 2009).

Parvalbumin-Expressing Interneurons

Parvalbumin-positive interneurons are widespread throughout the brain and can be subdivided by their morphological characteristics, as well as their laminar and anatomical locations (Freund & Buzsaki, 1996; Hof et al., 1999). PV protein is primarily expressed in three distinct GABAergic interneuron subtypes: basket cells, chandelier cells (in neocortex; referred to as bistratified cells in HPC), and O-LM cells (in stratum oriens in HPC) (Freund & Buzsaki, 1996; Klausberger, 2009). Chandelier cells are primarily located in cortical layers II-VI, as well as within the hippocampus (though here they are often referred to as axo-axonic or bistratified cells), and have axonal terminations exclusively on the axon

initial segment of excitatory pyramidal neurons (Markram et al, 2004). Though there are relatively few of this cell type compared to other inhibitory interneurons, these are thought to be very powerful with regard to their ability to regulate neural transmission in the brain (Woodruff et al., 2010). Due to their inhibition directly onto the axon initial segment, chandelier cells also play an important role in regulating pyramidal cell excitatory output (Lewis et al., 2004); specifically, the ability for a single chandelier cell to target large populations of pyramidal neurons allows for wide-spread synchronization of excitatory transmission (Cobb et al., 1995).

Basket cells, which are located within the cortex, hippocampus, and cerebellum (Seto-Ohshima et al., 1989), synapse onto pyramidal cell somata and proximal dendrites and are primarily located within the pyramidal and granule cell layers in the hippocampus (Nitsch et al., 1990; Freund & Buzsaki, 1996). Like chandelier cells, a single parvalbumin-containing basket cell is able to make a large number of connections onto local pyramidal neurons, as well as synapsing with other parvalbumin-containing interneurons (Sik et al, 1995), thereby giving these interneurons the ability to strongly modulate postsynaptic excitatory output (Aradi et al., 2002). PV expressing cells are also located outside of the pyramidal and granule cell layers within stratum oriens, where these PV-expressing interneurons are referred to as O-LM cells, as well as in the hilus (Sloviter, 2004).

Within the CA1 region of the hippocampal formation, basket cells and chandelier cells make up approximately 75% of the total interneuron population (Szilagyi et al., 2011). The anatomical distribution of parvalbumin-positive

inhibitory interneurons within each sub-area of the hippocampus (CA1, CA3, dentate gyrus) appears to have consistent laminar profiles across the dorso-ventral axis, as well as no dorso-ventral differences in expression within each hippocampal area, at least in the rat (Kosaka et al, 1987; Nomura et al, 1997a; 1997b).

Treatment and Age-Related Changes in Parvalbumin Expression

Of particular relevance to the selective decrease in PV expressing interneurons within schizophrenic postmortem tissue, have been studies employing acute and/or chronic NMDA-antagonist drug regimens in rodents (e.g. Kittelberger et al., 2012). While such studies have generally reported a decrease in PV as a result of NMDA-antagonist administration (e.g. Romon et al., 2011; Wang et al., 2008; Braun et al., 2007; Keilhoff et al., 2004), there have been some studies indicating no change in PV-immunoreactivity following drug administration (e.g. Benneyworth et al., 2011; Zhang et al., 2008). What is interesting regarding these conflicting findings is that the studies do not control for age as a possible factor influencing PV expression. Rather, many of these studies seem to assume stability in PV expression over the lifespan. A major rationale for the current studies was to examine age-related differences in PV expression so as to better understand the existing literature and inform future studies on the effects of chronic NMDA-antagonist treatments.

Age-related changes in parvalbumin expression have been found in cortical (Ouda et al., 2008), as well as subcortical areas including the

hippocampus (Vela et al., 2003). However, there have been very few studies offering conclusive findings outlining age-dependent changes in hippocampal PV expression in a systematic way. In fact, to our knowledge, only a few studies have been conducted seeking to directly compare PV expression in young and aged rats in an attempt to characterize calcium-binding protein distribution as a function of age (e.g. Shetty & Turner, 1998; Vela et al., 2003; Stanley et al., 2012). Though there have been few studies, Shetty and Turner (1998) describe significant decreases in PV immunoreactivity within HPC CA1, CA3, and hilus between 5- and 24-month old rats, and these findings are in agreement with the limited work conducted across portions of the rat lifespan (Vela et al., 2003; Stanley et al., 2012).

The lack of systematically conducted normative baseline data regarding PV distribution across the lifespan is concerning, especially considering that rodent studies examining PV expression rarely account for the age of the animals used. This is evident in a study conducted by Nomura and colleagues (1997a) reporting no changes in hippocampal PV expression. Though they looked both between hippocampal regions and across the septotemporal axis, their data was exclusively collected from 5 week old rats, and therefore should not be generalized as a stable property of PV distribution across all age groups. Furthermore, the lack of baseline data makes it difficult to make conclusions based on data derived from experiments utilizing drug manipulations, as PV immunoreactivity may be dynamically altered based on a variety of endogenous and exogenous factors including, but certainly not limited to, age and experience.

The current study aimed to provide the first systematic characterization of PV-expression within the hippocampus, as well as within somatosensory and limbic cortices. In order to address the lack of data on normative PV distribution we set out to determine the following things with regard to PV immunoreactivity in young, adult, and aged rats: 1) whether PV expression differs along the septotemporal axis of the hippocampus; 2) whether or not PV exhibits an age-dependent profile of distribution and to determine whether our findings were consistent with the few studies which have also looked at age as a key factor; and 3) whether PV expression varies as a function of hippocampal sub-region (CA1, CA3, and DG), as well as cortical region (somatosensory and limbic cortices). The current study provides key normative data to inform future studies employing experimental manipulations, and allows for a more dynamic view of PV as a neural marker for pathology.

Chapter Two: Manuscript

Abstract

Parvalbumin (PV)-expressing inhibitory interneurons are the most numerous GABAergic interneuron type in the neocortex and hippocampus (HPC). Selective reductions in PV interneurons are observed in cortical and limbic regions in postmortem human schizophrenic brain tissue. Reductions in PV expression are also observed in rodents following acute and/or chronic treatment with NMDA-antagonists (e.g. ketamine). There are inconsistencies in the literature regarding age-related changes in PV expression in the HPC; some stating age-related decreases, no observed changes, and even increases in PV-immunoreactivity. These inconsistencies make it difficult to qualitatively and quantitatively evaluate changes observed in rodent models following NMDA-antagonist treatment. In order to provide a better baseline understanding of putative developmental changes in normal PV expression, the current study quantified 1) age-related changes in HPC and cortical PV expression in rats; 2) whether PV varied along the septotemporal axis of the HPC; and 3) regional changes in HPC PV expression. PV immunohistochemical methods were used to examine PV expression in hippocampal and adjacent neocortical regions in 1 and 6 month old naïve, as well as 24 month old behaviorally trained (radial water maze training), male Sprague-Dawley rats. Our main findings include: 1) specific regional differences (CA1, CA3, DG) across the septotemporal axis that varied by developmental age; 2) age-related decreases in PV expression in CA1 and DG (50-68% and 88-89% decrease from 1 – 6 months, respectively) in naïve rats;

and 3) A prominent increase in PV expression in the 24 month old rats that may be related to age or behavioral experience. These findings demonstrate developmental differences in the regional expression of PV, as well as potential behavioral and experiential increases in PV expression. Importantly, these baseline data provide key information for future work examining age-related differences in PV as a consequence of acute or chronic NMDA-antagonist treatment.

Introduction

A consistent finding in post-mortem schizophrenic brain tissue has been a selective reduction in a subset of inhibitory GABAergic interneurons containing the calcium-binding protein parvalbumin (PV) (Beasley & Reynolds, 1997; Volk et al., 2012; Lewis et al., 2012). PV-containing interneurons provide critical inhibition onto the soma and proximal dendrites of excitatory pyramidal neurons (Zuschratter et al., 1985; Kawaguchi et al., 1987; Katsumaru et al., 1988; Williams et al., 1992), which allows them to play a key role in determining the excitatory output of their postsynaptic targets (Lewis et al., 2004). This role likely contributes to the ability for involved neuronal circuits to accurately relay relevant information and support neural circuit processes key to cognitive function. Because of their importance in neocortical and hippocampal neuronal circuitry, the decrease in PV interneurons seen in schizophrenic brains likely contributes to the cognitive dysfunctions observed in schizophrenia.

Research of the underlying etiology of schizophrenia is difficult, as post-mortem human tissue can be confounded by countless variables (e.g., medication, co-morbid illnesses, drug use, etc.). Therefore, rodent models have been integral in enhancing our understanding of the proposed neural mechanisms underlying the disorder. A multitude of rodent models exist for the study of schizophrenia, ranging from genetic knockout mice (Chen et al., 2006), social isolation (Stevens et al., 1997), post-natal (Wedzony et al., 2008) or adult drug administration (Egerton et al., 2005; Chrobak et al., 2008), and even those employing a “two-hit” model consisting of two separate insults over the lifespan

(e.g., Bayer et al., 1999; Corriveau & Glenn, 2012). Of particular interest are rodent models exhibiting NMDA-antagonist induced selective reductions in PV expression in key brain regions such as prefrontal cortex and hippocampus (Keilhoff et al., 2004; Morris et al., 2005; Abekawa et al., 2007; Zhang et al., 2008). However, these findings often lack consistency, with some researchers indicating no changes in PV expression (Benneyworth et al., 2011), and other reports indicating exactly the opposite findings of one another (see Benneyworth et al., 2006; table 3). Alarming, there have been few studies providing consistent information regarding what could be considered 'normal' with regard to PV distribution within rodent brains, and whether the changes we are seeing are truly representative of human schizophrenic pathology.

In order to better understand the normative distribution of PV interneurons throughout the brain in rodents, and to establish a baseline of PV distribution consistent with past findings, we conducted a series of immunohistological studies in rats to visualize PV expression in cortical and hippocampal areas. We also quantified PV expression across the rat lifespan (specifically in post-weaning, young adult, and aged cohorts) within the brain regions of interest, as there are mixed reports indicating a decline in PV expression in some (Lolova & Davidoff, 1992; Miettinen et al., 1993), but not all (e.g., Vela et al., 2003; Potier et al., 2006), age-related PV studies. The current study also looked across the septotemporal axis of the hippocampus across aging, as there have been no studies, to our knowledge, that have systematically examined changes in PV expression in such a way. Here, we provide compelling evidence for an age-

dependent reduction of PV expression across hippocampal areas, as well as age-dependent septotemporal variation in PV expression. Furthermore, we provide evidence suggesting a role for behavioral enrichment in increasing PV expression within the rodent brain. Overall, our findings support a dynamic role of PV expression across the rat lifespan, particularly within the hippocampus and, to a lesser degree, within adjacent limbic cortex and neocortical regions.

Materials and Methods

Animals

Twenty-six male Sprague Dawley rats (Charles River Laboratories, Wilmington, MA) spanning three different ages were used: adolescent (approximate age P30; referred to as 1 month old; $n = 8$), young adult (approximate age P120-180, referred to as 6 month old; $n = 8$), and aged (24 months old, $n = 8$). Young adult and aged rats were single-housed, while adolescent rats were housed in groups of four, in clear polycarbonate caging. Normal husbandry procedures were followed for all rats within a temperature and humidity controlled vivarium, with a 12-hour light/dark cycle (lights on at 0800h), with access to food and water *ad libitum*. Additional animals were used to optimize Parvalbumin (PV) antibody concentration prior to immunohistochemical processing of experimental tissue ($n = 2$).

Young (1 month) and adult (6 month) rats were experimentally naïve, while aged (24 month) rats had a history of behavioral testing (continuous radial

water maze experience from approx. P60 – 1.5 years of age; details of training can be found in Chrobak et al., 2008) and pharmacological testing (ketamine and cannabinoid antagonist treatments from 1-1.5 years of age). Additionally, aged rats had restricted food intake strictly during behavioral testing. However, all behavioral and pharmacological manipulations were completed minimally four months prior to sacrifice/brain collection. All housing and experimental procedures were in accordance with, and approved by, the University of Connecticut Institutional Animal Care and Use Committee.

Immunohistochemistry

All rats were anesthetized with Euthasol (pentobarbital sodium solution; 1mL via intraperitoneal injection) and transcardially perfused first with ice-cold .9% physiological saline solution (approximately 200mL for 5 minutes), followed by ice-cold 3.7% paraformaldehyde solution (approximately 200mL for 5 min). Brains were removed and stored in 3.7% paraformaldehyde solution prior to being cryoprotected in a 30% sucrose solution for 48 hours before being sliced into 60µm coronal sections on a cryostat. Sections were taken from approximately -2.5mm through -6.0mm relative to Bregma (Paxinos & Watson, 1997; see Figure 1) that included distinct septotemporal areas of the HPC as well as adjacent cortical areas (retrosplenial and somatosensory), and then stored in phosphate-buffered saline (PBS)-containing well plates.

One series of tissue slices (each sample roughly separated by 240µm) per brain were selected to examine Parvalbumin (PV) immunoreactivity ($n=24$

series). Free-floating sections were initially blocked in 5% normal goat serum (Jackson Immuno), 0.1% triton-X, and PBS solution for 1h, followed by 3 five-minute rinses in PBS. Sections were then transferred into primary antibody solution (1:4000 anti-rabbit parvalbumin antibody (Calbiochem), 0.1% triton-X, and PBS – antibody concentration determined previously using aged rat brains ($n=2$) (see Figure 2)) for a 24h incubation period at room temperature. Sections were then rinsed 3 times at 5 minutes, and transferred into secondary solution (HRP labeled polymer anti-rabbit, DAKO) for a 2h room-temperature incubation period. Following a final set of 3 five-minute rinses in PBS, sections were transferred into a diaminobenzidine (DAB) chromogen solution (DAKO; Carpinteria, CA) for ten minutes to develop the stain. Sections were mounted and dried on glass slides, then cover-slipped for microscopy using DPX. Adjacent sections were processed for Nissl body staining using thionin to assist with identification of brain regions.

Data Analysis

A Nikon Eclipse E600 (Melville, NY) upright microscope equipped with an Insight SPOT digital camera (Diagnostic Instruments, Inc.) was used to examine and photograph sections with the assistance of SPOT software (SPOT Imaging Solutions; Sterling Heights, MI). Sections containing the major hippocampal subdivisions (CA1, CA3, and DG) were selected for analysis along the septotemporal axis within -2.45mm through -3.9mm (septal regions) and within -5mm through -6mm (midseptotemporal and temporal regions) relative to Bregma

(Steullet et al., 2010; Nomura et al., 1997a); see Figure 1 for section selection. Additionally, photomicrographs were taken of retrosplenial and somatosensory (barrel field) cortices, sampled from sections containing septal hippocampus, and were digitally catalogued for later analysis (see Figure 10A). Hippocampal sub-regions and retrosplenial cortex were photographed at 10x magnifications, while neocortical areas were photographed at 4x magnification to ensure inclusion and visualization of all cortical layers.

Digital photographs were analyzed using ImageJ software (NIH), and the number of PV immunoreactive interneurons was quantified using macros written to automate particle counting within each specified region. The sizes of the analyzed regions were as follows: CA1/CA3/ DG, 800 μ m x 500 μ m; retrosplenial cortex, 800 μ m x 800 μ m; somatosensory, 1600 μ m x 1000 μ m. Sections were chosen for analysis based on their anatomical location; representative photomicrographs containing septal CA1, CA3, and DG, were taken from levels corresponding to -2.45mm to -3.9mm relative to Bregma (Figure 3A); midseptotemporal and temporal CA1, CA3, and DG were taken from levels corresponding to -5.0mm to -6.0mm relative to Bregma (Figure 3B) (Paxinos & Watson, 1997). The average PV immunoreactivity, as counted using ImageJ macro parameters (see Figure 4), for each counted region was averaged for each age group to determine whether there were differences across hippocampal regions and/or across the long axis, in addition to possible differences within neocortical regions. Regional and areal differences in the distribution of PV+

interneurons were assessed using one-way Analysis of Variance (ANOVA) followed by post-hoc t-tests (Kirk, 2012).

Results

Summary of Results

Overall, data indicated an age (1-, 6-, and 24-month), regional (cortical field, hippocampus), septotemporal (septal, midseptotemporal, temporal), and hippocampal region (CA1, CA3, DG) variation in the distribution of PV-expressing interneurons. First, we will describe the differences in PV expression across the septotemporal axis within the hippocampal subdivisions when comparing 1- and 6-month old rat brain tissue. In 6 month old rats, there was a significant difference in PV expression from septal to temporal CA1 and DG (50- 80% decrease in expression from septal to temporal regions). Second, we describe the age-dependent differences in PV expression observed within specific hippocampal subdivisions. Generally, there were significant decreases in the number of PV expressing neurons in the CA1 and DG regions of 6-month, compared to 1-month, old rats without concomitant changes within the CA3 sub-region. No differences were observed in the two neocortical fields from the same septal coronal sections (barrel field and retrosplenial cortices). Third, we describe the regional (CA1, CA3, DG) differences in PV expression, where there is, generally, a higher density of PV+ interneurons observed in CA1 sub-regions in contrast to CA3 or DG. Lastly, we report a striking increase in PV expression in all regions of interest (cortical areas and hippocampal regions) in the 24-month

old age group, compared to 6-month old rats. Notably, this effect is confounded by the extensive behavioral training that the 24-month old rats received. Ongoing experiments are being conducted to determine whether these differences are consequent of age and/or experience.

Age-related septotemporal variation in HPC parvalbumin expression

There was no difference in the distribution of PV expressing neurons across the septotemporal axis of CA1, CA3, or the DG in 1 month and 24 month old rats (see Figure 5). In contrast, 6-month old rats exhibit a significant effect of septotemporal location in CA1 and DG. Specifically, septal CA1 in 6 month old rats had significantly greater PV expression than both midseptotemporal and temporal CA1 ($F(2, 14) = 21.4, p < .001$; see Figure 5A). Similarly, septal DG in 6-month old rats had significantly greater PV expression than the midseptotemporal region ($F(1, 7) = 6.5, p < .05$; see Figure 5B). In contrast, there were no septotemporal variations in region CA3 of the hippocampus ($F(2, 14) = 3.3, p = .07$; see Figure 5C). For representative photomicrographs of CA1, CA3, and the DG, see Figures 6, 7, and 8 (respectively); descriptive statistics can be seen in Table 1.

1- to 6-months: Age-related difference in regions CA1 and DG that increases with difference from septal pole

Overall, there was a similar expression of PV in region CA3 in 1 and 6 month old rats across all septotemporal levels, however, there was less PV in

more temporal levels in both CA1 and the DG. Specifically, there was an age-dependent decrease in PV expression within CA1, such that 1-month old rats had significantly higher PV expression than 6-month old rats in midseptotemporal ($t(14) = 3.3, p < .01$) and temporal levels ($t(13) = 3.8, p < .01$), though there was no difference between these groups in septal CA1 ($p = .24$); see Figure 9A. There was also an age-dependent decline in PV expression in both septal DG ($t(14) = 7.1, p < .001$) and midseptotemporal DG ($t(12) = 2.4, p < .05$), see Figure 9B. However, no age-related differences in PV expression were observed in septal ($p = .11$), midseptotemporal ($p = .66$), or temporal ($p = .47$) CA3 (Figure 9C). These data can be visualized in representative photomicrographs in Figures 6, 7, and 8. No significant age-related effects were seen in retrosplenial ($p = .52$; Figure 10B) or barrel field ($p = .28$; Figure 10C) cortices between 1- and 6-month old naïve rats (Representative photomicrographs in Figures 11 and 12). Descriptive statistics can be seen in Table 1.

CA1 regions have increased PV expression compared to other HPC regions across all age groups

Significant regional differences in PV expression were seen in septal and midseptotemporal, but not temporal, hippocampal regions across all age groups (1-, 6-, and 24-months). 1-month old rats have significantly higher PV expression in septal CA1 compared to septal CA3 ($p < .01$) and DG ($p < .001$). Midseptotemporal sections follow the same pattern, with CA1 having significantly more PV expression than CA3 ($p < .001$) and DG ($p < .001$). 6-month old rats had significantly higher PV expression in septal CA1 in comparison to septal CA3

($p < .001$) and DG ($p < .001$), in addition to having significantly higher PV expression in midseptotemporal CA1 compared to both midseptotemporal CA3 ($p < .05$) and DG ($p < .01$). In 24-month old extensively trained rats, septal CA1 had significantly higher PV expression than septal CA3 ($p < .01$) and DG ($p < .001$). Interestingly, in midseptotemporal regions, CA1 and CA3 were not significantly different from one another ($p = .25$), but both had higher PV expression than midseptotemporal DG ($p < .01$, $p < .05$; respectively). These data can be seen in Figure 13, and descriptive statistics can be seen in Table 1.

Increased parvalbumin expression in 24-month old extensively trained rats across all hippocampal and cortical regions examined

A striking increase in PV expression was observed in 24-month old extensively trained rats relative to 1- and 6-month old naïve rats within all hippocampal and cortical regions quantified. There was significantly more PV expression in both retrosplenial and barrel field cortices in 24- compared to 1- ($p < .05$, $p < .001$; respectively) and 6- ($p < .05$, $p < .001$; respectively) month old rats (Figure 10). 24-month old rats had significantly higher PV expression than 6-month old rats in all hippocampal regions observed: Septal CA1 ($p < .05$), CA3 ($p < .05$), and DG ($p < .001$); midseptotemporal CA1 ($p < .01$), CA3 ($p < .01$), and DG ($p < .05$); and temporal CA1 ($p < .001$) and CA3 ($p < .01$). 24-month old rats had significantly higher PV expression than 1-month old rats in midseptotemporal ($p < .05$) and temporal ($p = .01$) CA3, but these two age-groups had statistically equivalent PV expression in all other hippocampal regions observed (p 's $> .1$). These data can be seen more clearly in Figure 14, and Table 1.

Discussion

Understanding the normative distribution of a target cellular population - in this case PV expressing GABAergic interneurons - is integral to the interpretation and understanding of data derived from experimental manipulations (e.g., pharmacological, trauma, etc.) in rodent models of psychiatric disorders. Currently, there is a significant lack of normative baseline knowledge of PV expression across the hippocampal septotemporal axis as a function of aging. The current data decreases the paucity of data in this regard and should contribute to future evaluations of PV loss consequent to experimental manipulations. Importantly, we provide compelling evidence of an age-dependent decline in PV expression that varies with respect to hippocampal region (CA1, CA3, and the DG), as well as across the septotemporal axis. Furthermore, we present evidence that PV expression may be more dynamic than previously thought and that extensive behavioral training may actually increase PV expression within the rat brain.

Septotemporal differences in hippocampal PV expression are age-dependent

It is well-known that the hippocampus is integral for cognitive processing and memory formation within the brain (Milner et al., 1998). The hippocampus can be sub-divided along its areal or longitudinal axis into: septal, midseptotemporal, and temporal levels (Amaral & Witter, 1989). These septotemporal regions are also associated with different cognitive, emotional,

and behavioral information processing. Generally speaking, septal regions of the hippocampus receive spatial information in addition to sensory information from the neocortex, while temporal hippocampus favors emotionally salient information processing, due to its increased amygdala input (Bannerman et al., 2004.) Because of these distinct differences in information processing within this structure, it follows that there would also be anatomical markers that vary parallel to functional variation across the septotemporal axis (Gusev et al., 2005; Jinno, 2011).

In the current study, we found clear septotemporal variations in PV immunoreactivity within the hippocampal formation of 6-month old behaviorally naïve rats. Specifically, there was a significant difference in PV expression from septal to midseptotemporal CA1 and DG, with septal areas exhibiting significantly more PV expression than midseptotemporal areas. Conversely, there were no differences in expression between midseptotemporal and temporal CA1 locations. However, this pattern was not observed in the CA3 region of 6-month old rats, nor were there any septotemporal differences in any region for 1-month old naïve and 24-month old behaviorally trained groups. The lack of septotemporal difference within the CA3 region in 6-month old rats suggests the possibility that PV expressing neurons in this region may be more resilient to aging, while CA1 and DG regions may have increased susceptibility to the adverse effects of aging. However, these findings may also indicate a dynamic interaction between aging and septotemporal location within the hippocampus as

well. To our knowledge, this is the first systematic characterization of septotemporal variation in PV expression across distinct age groups.

Previous findings from Nomura and colleagues (1997a) indicate no significant septotemporal variation in PV expression; however, their work was conducted in 5 week old rats (approx. 1 month of age) and therefore is in agreement with the lack of septotemporal effect seen in the 1-month old naïve rats in the current study. Conversely, in mouse brains processed to visualize parvalbumin-positive interneurons, research has shown significant dorsal-ventral differences in parvalbumin expression, particularly within stratum pyramidale (Jinno & Kosaka, 2006). More specifically, Jinno and Kosaka (2006) provide evidence of elevated levels of PV expression within CA1 compared to CA3 and DG, which is in agreement with evidence presented in the current study for 6 month old rats.

It is difficult to interpret the lack of septotemporal variation in PV expression within the hippocampus of 24-month old behaviorally trained rats. This is primarily due to the fact that this was the only group that received extensive behavioral training and it is possible that the training influenced the amount of PV expression throughout the brain, including those areas examined presently. Conversely, it is also possible that there is an age-dependent decrease, then subsequent increase, in PV expression. Either way, these results shed light on the dynamic properties of PV protein expression across the hippocampus.

Age-dependent decline in PV expression within the rat hippocampus

Our data provide support for a growing body of evidence describing an age-dependent decline in PV expression within the hippocampus. Work by Nomura and colleagues (1997a) indicate a high degree of PV expression within young (5 week old) rats that is in line with our findings in 1 month old rats of high levels of PV expression coinciding with a lack of septotemporal gradient. However, we observed a significant decline in PV expression as a result of age in CA1 and DG from 1-month to 6-month experimentally naïve rats, suggesting that expression may be developmentally dynamic. These findings are in agreement with previous studies reporting age-related decreases in PV expression within the hippocampus (Kuruba et al., 2011; Lolova & Davidoff, 1992). Importantly, our data argues against previous research indicating an absence of age-related decline (e.g. Kishimoto et al., 1998) or the presence of age-related increases (Choi et al., 2010) in hippocampal and cortical PV expression. Furthermore, the lack of age-dependent effects of PV immunoreactivity in CA3 suggest the possibility that age-related decline in PV expression may be regionally dependent within the rat.

Data from 24-month old rats was not included in the aging analysis because of the anomalous findings which may be a result of the extensive behavioral training these rats received over their lifetime. Generally, 24-month extensively trained rats exhibited equal (if not more) PV expression in comparison to 1-month old rats across all hippocampal sub-regions (Figure 14),

as well as cortical regions (Figure 10). In addition, 24-month old rats had significantly higher PV expression in all hippocampal sub-regions and cortical regions (with the exception of no significant difference in septal CA3, though a clear trend was present; Figure 14). While we provide evidence against an age-related loss of PV immunoreactivity in 24 month old rats, we cannot claim that this is an expected and normative finding in aged rat hippocampus. In fact, in light of the age-related decline in PV expression from 1- to 6-month old naïve rats, we suggest that these findings are an artifact of the extensive behavioral training this group received. Furthermore, we hypothesize that, if these animals had not received a lifetime of behavioral training, their PV expression profile would be that seen currently in the 6-month old naïve rats, if not further decreased, providing strong evidence for PV decrease across the lifespan (see Figure 15 for a proposed timeline of PV expression).

Possible role of extensive training/enrichment in increasing overall PV expression

The prominently high levels of PV expression within all regions of interest in 24-month old extensively trained rats in the present study were surprising and unexpected. While we initially set out to characterize possible age- and regionally-dependent differences in PV expression, we observed consistently elevated expression both within the hippocampal regions (Figure 14), as well as the cortical regions (Figure 10) examined. It is unlikely that these findings are a result of an age-dependent shift in PV immunoreactivity, but rather suggest a

dynamic interaction between the expression of this particular calcium-binding protein within the brain and the behavioral experiences of the animal.

Experience-related changes in the expression of neuronal and cellular markers (e.g., BrdU, NeuN, BDNF, apoptotic markers, etc.) are a well-established phenomenon in the literature (van Praag et al., 2000). A multitude of experiences have the capability of altering brain structure and function, including (but certainly not limited to): maze training (Van der Borght et al., 2005), social interaction (Woolley & Doupe, 2008); physical stimulation (e.g. stimulation of whiskers; Knott et al., 2002); sexual experience (Leuner et al., 2010); stress (Zoladz et al., 2012); and, a topic of extensive research, environmental enrichment (Harati et al., 2011). There are few studies which have examined the effects of enrichment specifically with regard to PV, but those that have been reported indicate enrichment and/or deprivation-related alterations within sensory cortices (Jiao et al., 2006), striatum (De Bartolo et al., 2011), and hippocampus (Iuvone et al., 1996). The findings by Iuvone and colleagues (1996) provide evidence of experience-related increases in PV expression, specifically within the hippocampal regions CA1, CA3, and DG, which is consistent with our findings in 24-month old rats, and suggests that this finding is likely an artifact of enrichment, and not an age-related effect.

General Discussion

Overall, we present evidence in support of age-related decreases in PV-expressing GABAergic interneurons within CA1 and DG regions of the

hippocampus from 1- to 6-month old naïve rats, as well as regionally-specific expression in CA1 compared to CA3 and DG across three distinct age groups. We also provide compelling evidence for a dynamic interaction between PV protein expression within the hippocampus and neocortical regions and life-long enrichment in the form of extensive behavioral training. The present findings provide a framework to inform future studies. Specifically, the regional and age-related differences seen here as a result of age can allow for future experiments seeking to understand the effects of pharmaceutical agents (e.g. NMDA-antagonists), such that they can ensure proper controls, and therefore provide more compelling evidence for differences in neuronal markers. Additionally, the results seen in extensively trained animals open up a new avenue of research, and could possibly inform preventative studies of psychopathologies. Though preliminary in nature, these data provide straightforward evidence for age- and experientially-dependent influences on the expression of PV in rat hippocampus and nearby sensory cortices.

Chapter Three: Discussion

The current study provides support for past research indicating an age-dependent decrease in PV expression within the hippocampus of rats (e.g. Vela et al., 2003). Furthermore, it provides new and compelling evidence for an age-dependent decline in PV expression across the septotemporal axis, which, to our knowledge, has not been previously reported. We found evidence indicating elevated PV expression within the hippocampal CA1 sub-region, compared to CA3 and DG regions, across all age-groups, and a general lack of differences within the CA3 sub-region across age and septotemporal location. Of particular interest, we describe possible experientially-dependent effects on PV expression, indicating that PV has dynamic expression, and can likely be differentially affected by a variety of environmental and biological phenomena. The current study is the first of its kind in systematically characterizing PV expression not only across the septotemporal axis of the hippocampus, but also across diverse age groups in rats, and provides critical baseline data describing PV expression within key brain regions implicated in psychopathology.

Contributions of the current work to the literature: Age-related changes in PV expression

Previous work detailing the distribution of PV expression within the rodent hippocampus and other cortical areas has been important to our understanding of their role in interneuronal circuits. For instance, Nomura and colleagues

(1997a) characterized PV expression across the septotemporal axis of the hippocampus, and indicated no differences in PV expression based on septotemporal level. These findings perpetuate the notion that PV expression is a static marker in GABAergic interneurons within the normal, untreated brain. However, this study lacks generalizability across age groups, as their analyses were only performed in young rats (5 weeks of age). This is problematic for a variety of reasons, but particularly so because many experimental studies (e.g. those using NMDA-antagonist treatments) do not control for age, and baseline data for PV expression across age-groups is needed for comparison and accurate interpretation of results.

An overwhelming degree of inconsistency is seen in the reports of PV immunoreactivity following acute and/or chronic administration of NMDA-antagonist drugs (e.g. ketamine; MK-801). We believe that many of these inconsistencies are likely a result of a failure to control for age in the rodent models utilized, and that their questions may be better answered if they were aware of the dynamic changes seen in PV expression as a consequence of age. For example, studies that have seen no significant changes in PV expression, especially in the hippocampus, following NMDA-antagonist administration regimens have used rodents aged PD7 (Wang et al., 2008), 1 month of age (Zhang et al., 2008), PD 20 & 40 (Stuellet et al., 2010); and 3 months of age (Benneyworth et al., 2011). Note, that all of these studies, which have found no significant impact of NMDA-antagonism on PV expression, have not only used a wide variety of ages across studies, but all of these animals have generally been

3 months of age or less. According to our findings, 1 month old rats appear to have very high expression of PV across all regions studied, and this expression is likely robust and resistant to change in young animals.

Conversely, studies conducted administering NMDA-antagonist drugs in rats 4 months of age and older, observed significant decreases in PV expression within the hippocampal formation (Stuellet et al., 2010; Keilhoff et al., 2004), as well as within the prefrontal cortex (Romon et al., 2011). This disparity is compelling, as it suggests that there may be an age-dependent response to NMDA-antagonist drugs, making adult rodents more susceptible to PV loss as a result of administration of these substances. This could perhaps indicate a decreased stability in the expression of PV in adulthood or, more simply, that PV is a dynamic protein that has the ability to constantly up- or down-regulate its expression with the age and needs of the animal. To make this argument more compelling, studies should be conducted in aged rats (1+ years) in these NMDA-antagonist models to determine whether a similar pattern to adult PV loss is seen, or if perhaps PV decreases are limited to a window of susceptibility, much like what is seen for schizophrenic onset in humans (e.g. Gogtay et al., 2011; Castle et al., 1998). Overall, the current work sheds light on the need for considering age in rodent models of psychiatric disorders to account for the dynamic expression of neuronal markers (e.g. PV) throughout the lifespan.

Contributions of the current work to the literature: Experience-related changes in PV expression

Here we describe a new and intriguing phenomenon of experience-induced increases in PV expression in aged (24-month old) rats. To our knowledge, this is the first report of increased PV in aged rats as a result of extensive life-long behavioral training. While previous studies have observed increases in PV as a result of experience, these results are limited, and generally encompass young animals. For example, reports of increased PV expression within the brain have been attributed to enrichment-related events such as: neonatal auditory stimulation (Chaudhury et al., 2008); increased neural activity in visual cortex (Patz et al., 2004); univibrissa rearing (Nowicka et al., 2009); and adolescent exercise (Gomes da Silva et al., 2010), among others. Furthermore, sensory deprivation in young rodents has been shown to down-regulate PV expression (e.g. Jiao et al. 2006).

Interestingly, a study by Siucinska and Kossut (2006) reports no difference in PV expression in the barrel cortex as a result of short-term sensory learning in adult mice. This is contradictory to our findings of experience-related increases in PV expression within the barrel cortex in 24-month old rats. However, Siucinska and Kossut looked at the effects of short-term sensory learning, while our effect is driven by extensive, life-long behavioral training and experience. Though younger animals seem to have a greater benefit of experience on PV expression, it may be the case that adult and aged animals need more extensive experience in order for it to manifest with increased PV expression. Our data provide

descriptive and quantitative evidence for an up-regulation of PV expression following extensive behavioral training, and can be taken as evidence for the dynamic nature of PV expression across the lifespan, and when faced with different short- or long-term experiences.

Future Directions

While preliminary in nature, the current study has provided us with novel and informative data to move forward with on-going studies in our lab to understand anatomical consequences of chronic NMDA antagonist treatment. Furthermore, the baseline data we have collected through this study provide a starting point for more in-depth future studies to more thoroughly characterize the dynamic nature of PV expression across the lifespan. Of particular note, the data indicating a significant increase in PV expression in extensively trained aged rats has provided rationale for a new avenue of research describing enrichment-induced changes in calcium-binding protein expression.

Expanding upon observed age-dependent changes in PV expression

While the current study provides a starting point for more thorough research into age-related differences in PV expression throughout the rodent brain, there is still much to be done. To our knowledge, there are not studies which have systematically explored PV expression across multiple brain regions as a function of age, and particularly those that look at rats over 1.5 years of age

in comparison to postnatal and adult rats. Such characterization would not only be beneficial to the overall understanding of PV distribution within the rodent brain, but would provide reliable baseline data from which to base future studies aiming to observe PV expression as a meaningful marker of pathology.

In an attempt to further understand the effect of age on PV expression, it would be useful to conduct a series of studies looking at multiple ages of rats including: postnatal pups, young (e.g. 1-3months), young adults (e.g. 6 months), adult (e.g. 1 year) and aged (e.g. 2 years) in order to describe the dynamics of PV expression as a function of age. Furthermore, within each age group multiple brain regions should systematically examined including: hippocampal sub-regions (CA1, CA3, DG, subiculum), somatosensory cortices, limbic cortices, as well as other regions known to have high expression of PV (e.g. cerebellum; De Bartolo et al., 2011). Such an anatomical characterization of PV distribution would add depth to previous studies which have attempted to provide comprehensive understanding of PV distribution, but did not account for age and had a more narrow anatomical focus (e.g. Nomura et al., 1997a; 1997b).

Expanding upon observed enrichment-dependent changes in PV expression

We are especially interested in following up on the enrichment-induced changes observed in the 24-month old rats in the present study. As this is the first report of extensive training resulting in a substantial increase in PV expression (back to levels seen in 1-month old naïve rats and sometimes even

significantly higher), follow-up work is required to validate these results in a more controlled manner. Specifically, we propose to more closely examine the role of enrichment across the rat lifespan to determine 1) whether PV expression is, as is suggest in the present body of work, age-dependent in naïve rats; 2) whether septotemporal variation in PV is age-dependent, as our data imply; 3) whether behavioral experience (in the form of extensive radial water maze training) significantly increases PV expression throughout hippocampus and other areas of the brain (e.g. somatosensory and limbic cortices); and 4) whether an interaction exists between age and experience to differentially modify PV expression in the rat brain. We hypothesize that in naïve rats a significant age-dependent decrease in PV expression would be evident, and that rats receiving extensive training would have no age-dependent decrease in PV expression compared to their naïve counterparts; findings which would be consistent with the data presented in the current study.

Due to the wide use of decreased PV expression as a marker in animal models of schizophrenia (e.g. Kittelberger et al., 2012; Benneyworth et al., 2011), our lab is also interested in the effects of NMDA-antagonism (ketamine) on PV expression and memory as a function of age, as well as experience. Within the radial water maze paradigm (described in Chrobak et al., 2008), we suspect that ketamine-induced memory deficits in the radial water maze may be mirrored by a concomitant decrease in PV expression. Furthermore, since these animals will have had extensive training, and therefore increased PV expression, we believe that any ketamine-induced changes in expression would be clearly evident. In

fact, decreases in PV expression in ketamine-treated behaviorally trained rats are hypothesized to have similar, if not the same amount, of PV expression as their aged experimentally naïve counterparts. Such findings would provide compelling evidence of PV as a dynamic and fluctuating biological marker that has the ability to be influenced by biological, pharmacological, and environmental factors. Furthermore, these proposed findings would also provide validity to previous studies which have observed NMDA-antagonist induced decreases in PV expression (e.g. Kittelberger et al., 2012) and perhaps even offer explanatory evidence for why other studies failed to see a decrease as a result of NMDA-antagonist administration due to their sole use of young rats (e.g. Benneyworth et al., 2011).

General Conclusions

Overall, the current body of work has provided a solid foundation for future work describing age-related and septotemporal changes in PV expression in rodents. Additionally, it has opened up a new avenue of research into the dynamic properties of the calcium-binding protein, PV, and its modulation by behavioral training and enrichment. Though preliminary, our findings provide promising evidence for age- and experientially-dependent changes in PV expression with key brain regions associated with cognitive function. Importantly, the current findings of our study, and those we have proposed, will allow for better understanding, and interpretation, of future studies, especially with regard to modeling schizophrenic cognitive dysfunction.

References

- Abekawa T, Ito K, Nakagawa S, Koyama T (2007) Prenatal exposure to an NMDA receptor antagonist, MK-801 reduces density of parvalbumin-immunoreactive GABAergic neurons in the medial prefrontal cortex and enhances phencyclidine-induced hyperlocomotion but not behavioral sensitization to methamphetamine in postpubertal rats *Psychopharmacology* 192: 303-316.
- Amaral DG, Witter MP (1989) The three-dimensional organization of the hippocampal formation: A review of anatomical data *Neuroscience* 31(3): 571-591.
- Aradi I, Santhakumar V, Chen K, Soltesz I (2002) Postsynaptic effects of GABAergic synaptic diversity: Regulation of neuronal excitability by changes in IPSC variance *Neuropharmacology* 43(4): 511-522.
- Arif SH (2009) A Ca(2+)-binding protein with numerous roles and uses: parvalbumin in molecular biology and physiology *Bioessays* 31(4): 410-421.
- Bannerman DM, Rawins JN, McHugh SB, Deacon RM, Yee BK, Bast T, Zhang WN, Pothuisen HH, Feldon J (2004) Regional dissociations within the hippocampus-memory and anxiety *Neuroscience and Biobehavioral Reviews* 28(3): 273-283.
- Barch DM (2006) What can research on schizophrenia tell us about the cognitive neuroscience of working memory? *Neuroscience* 139: 73-84.
- Bayer TA, Falkai P, Maier W (1999) Genetic and non-genetic vulnerability factors in schizophrenia: the basis of the "two hit hypothesis" *Journal of Psychiatric Research* 33(6): 543-548.
- Beasley CL, Reynolds GP (1997) Parvalbumin-immunoreactive neurons are reduced in the prefrontal cortex of schizophrenics *Schizophrenia Research* 24: 349-355.
- Benes FM (2000) Emerging principles of altered neural circuitry in schizophrenia *Brain Research Reviews* 31(2-3): 251-269.
- Benneyworth MA, Roseman AS, Basu AC, Coyle JT (2011) Failure of NMDA receptor hypofunction to induce a pathological reduction in PV-positive GABAergic cell markers *Neuroscience Letters* 488: 267-271.

- Bergen SE, Petryshen TL (2012) Genome-wide association studies of schizophrenia: Does bigger lead to better results? *Current Opinion in Psychiatry* 25(2): 76-82.
- Bogerts B, Meertz E, Schönfeldt-Bausch R (1985) Basal ganglia and limbic system pathology in schizophrenia: A morphometric study of brain volume and shrinkage *Archives of General Psychiatry* 42(8): 784-791.
- Bourque F, van der Ven E, Fusar-Poli P, Malla A (2012) Immigration, social environment and onset of psychotic disorders *Current Pharmaceutical Design* 18(4): 518-256.
- Braun I, Genius J, Grunze H, Bender A, Möller HJ, Rujescu D (2007) Alterations of hippocampal and prefrontal GABAergic interneurons in an animal model of psychosis induced by NMDA receptor antagonism *Schizophrenia Research* 97: 254-263.
- Brown AS, Derkits EJ (2009) Prenatal infection and schizophrenia: A review of epidemiologic and translational studies *The American Journal of Psychiatry* 167(3): 261-280.
- Brown AS, Patterson PH (2011) Maternal infection and schizophrenia: Implications for prevention *Schizophrenia Bulletin* 37(2): 284-290.
- Buchanan RW, Freedman R, Javitt DC, Abi-Dargham A, Lieberman JA (2007) Recent advances in the development of novel pharmacological agents for the treatment of cognitive impairments in schizophrenia *Schizophrenia Bulletin* 33(5): 1120-1130.
- Burns JK (2004) An evolutionary theory of schizophrenia: cortical connectivity, metarepresentation, and the social brain *Behavioral and Brain Sciences* 27: 831-885.
- Buzsaki G, Chrobak JJ (1995) Temporal structure in spatially organized neuronal ensembles: A role for interneuronal networks *Current Opinions in Neurobiology* 5: 504-510.
- Camchong J, MacDonald AW, Bell C, Mueller BA, Lim KO (2011) Altered functional and anatomical connectivity in schizophrenia *Schizophrenia Bulletin* 37(3): 640-650.
- Carman J, Peuskens J, Vangeneugden A (1995) Risperidone in the treatment of negative symptoms of schizophrenia: A meta-analysis *International Clinical Psychopharmacology* 10(4): 207-213.

- Castle D, Sham P, Murray R (1998) Differences in distribution of ages of onset in males and females with schizophrenia *Schizophrenia Research* 33(3): 179-183.
- Chaudhury S, Nag TC, Wadhwa S (2008) Calbindin D-28k and parvalbumin expression in embryonic chick hippocampus is enhanced by prenatal auditory stimulation *Brain Research* 1191(29): 96-106.
- Chen J, Lipska BK, Weinberger DR (2006) Genetic mouse models of schizophrenia: From hypothesis-based to susceptibility gene-based models *Biological Psychiatry* 59(12): 1180-1188.
- Choi JH, Lee CH, Yoo KY, Hwang IK, Lee IS, Lee YL, Shin HC, Won MH (2010) Age-related changes in calbindin-D28k, parvalbumin, and calretinin immunoreactivity in the dog main olfactory bulb *Cellular and Molecular Neurobiology* 30: 1-12.
- Chrobak JJ, Hinman JR, Sabolek HR (2008) Revealing past memories: Proactive interference and ketamine-induced memory deficits *Journal of Neuroscience* 28(17): 4512-4520.
- Clarke MC, Tanskanen A, Huttunen M, Whittaker JC, Cannon M (2009) Evidence for an interaction between familial liability and prenatal exposure to infection in the causation of schizophrenia *The American Journal of Psychiatry* 166(9): 1025-1030.
- Cobb SR, Buhl, EH, Halasy K, Paulsen O, Somogyi P (1995). Synchronization of neuronal activity in hippocampus by individual GABAergic interneurons *Nature* 378: 75-78.
- Corriveau JA, Glenn MJ (2012) Postnatal choline levels mediate cognitive deficits in a rat model of schizophrenia *Pharmacology Biochemistry and Behavior* 103: 60-68.
- De Bartolo P, Gelfo F, Burello L, De Giorgio A, Petrosini L, Granato A (2011) Plastic changes in striatal fast-spiking interneurons following hemicerbellectomy and environmental enrichment *Cerebellum* 10(3): 624-632.
- Egerton A, Reid L, McKerchar CE, Morris BJ, Pratt JA (2005) Impairment in perceptual attentional set-shifting following PCP administration: A rodent model of set-shifting deficits in schizophrenia *Psychopharmacology* 179: 77-84.

- Freund TF, Buzsáki G (1996) Interneurons of the hippocampus *Hippocampus* 6(4): 347-470.
- Gogtay N, Vyas NS, Testa R, Wood SJ, Pantelis C (2011) Age of onset of schizophrenia: perspectives from structural neuroimaging studies *Schizophrenia Bulletin* 37(3): 504-513.
- Gold JM (2004) Cognitive deficits as treatment targets in schizophrenia *Schizophrenia Research* 72: 21-28.
- Gomes da Silva S, Dona F, Fernandes MJdS, Scorza FA, Cavalheiro EA, Arida RM (2010) Physical exercise during the adolescent period of life increases hippocampal parvalbumin expression *Brain and Development* 32(2): 137-142.
- Gonzalez-Burgos G, Lewis DA (2008) GABA neurons and the mechanisms of network oscillations: Implications for understanding cortical dysfunction in schizophrenia *Schizophrenia Bulletin* 34(5): 944-961.
- Greenwood TA, Swerdlow NR, Gur RE, Cadenhead KS, Calkins ME, Dobie DJ, et al. (2013) Genome-wide linkage analysis of 12 endophenotypes for schizophrenia from the consortium on the genetics of schizophrenia *The American Journal of Psychiatry In Advance*: 1-12.
- Gusev PA, Cui C, Alkon DL, Gubin AN (2005) Topography of Arc/Arg3.1 mRNA expression in the dorsal and ventral hippocampus induced by recent and remote spatial memory recall: dissociation of CA3 and CA1 activation *Journal of Neuroscience* 25(41): 9384-9397.
- Haijma SV, Van Haren N, Cahn W, Koolschijn PCMP, Hulshoff Pol HE, Kahn RS (2012) Brain volumes in schizophrenia: A meta-analysis in over 18,000 subjects *Schizophrenia Bulletin* [Epub ahead of print]: doi: 10.1093/schbul/sbs118.
- Harati H, Majchrzak M, Cosquer B, Galani R, Kelche C, Cassel JC, Barbelivien A (2011) Attention and memory in aged rats: Impact of lifelong environmental enrichment *Neurobiology of Aging* 32(4): 718-736.
- Harrison PJ (1999) The neuropathology of schizophrenia: A critical review of the data and their interpretation *Brain* 122(4): 593-624.
- Harrison PJ (2004) The hippocampus in schizophrenia: a review of the neuropathological evidence and its pathophysiological implications *Psychopharmacology* 174: 151-162.

- Hashimoto T, Volk DW, Egan SM, Mirnics K, Pierri JN, Sun Z, Sampson AR, Lewis DA (2003) Gene expression deficits in a subclass of GABA neurons in the prefrontal cortex of subjects with schizophrenia *The Journal of Neuroscience* 23(15): 6315-6326.
- Hof PR, Glezer II, Condé F, Flagg RA, Rubin MB, Nimchinsky EA, Vogt Weisenhorn DM (1999) Cellular distribution of the calcium-binding proteins parvalbumin, calbindin, and calretinin in the neocortex of mammals: phylogenetic and developmental patterns *Journal of Chemical Neuroanatomy* 16: 77-116.
- Holtzman CW, Shapiro DI, Trotman HD, Walker EF (2012) Stress and the prodromal phase of psychosis *Current Pharmaceutical Design* 18(4): 527-533.
- Honea R, Crow TJ, Passingham D, Mackay CE (2005) Regional deficits in brain volume in schizophrenia: A meta-analysis of voxel-based morphometry studies *American Journal of Psychiatry* 162(12): 2233-2245.
- Iuvone L, Geloso MC, Dell'Anna E (1996) Changes in open field behavior, spatial memory, and hippocampal parvalbumin immunoreactivity following enrichment in rats exposed to neonatal anoxia *Experimental Neurology* 139: 25-33.
- Jiao Y, Zhang C, Yanagawa Y, Sun QQ (2006) Major effects of sensory experiences on the neocortical inhibitory circuits *Journal of Neuroscience* 26(34): 8691-8701.
- Jinno S (2011) Topographic differences in adult neurogenesis in the mouse hippocampus: a stereology-based study using endogenous markers *Hippocampus* 21(5): 467-480.
- Jinno S, Kosaka T (2006) Cellular architecture of the mouse hippocampus: A quantitative aspect of chemically defined GABAergic neurons with stereology *Neuroscience Research* 56(3): 229-245.
- Katsumaru H, Kosaka T, Heizmann CW, Hama K (1988) Immunocytochemical study of GABAergic neurons containing the calcium-binding protein parvalbumin in the rat hippocampus *Experimental Brain Research* 72(2): 347-362.
- Kawaguchi Y, Katsumaru H, Kosaka T, Heizmann CW, Hama K (1987) Fast spiking cells in the rat hippocampus (CA1 region) contain the calcium-binding protein parvalbumin *Brain Research* 416(2): 369-374.

- Keilhoff G, Becker A, Grecksch G, Wolf G, Bernstein HG (2004) Repeated application of ketamine to rats induces changes in the hippocampal expression of parvalbumin, neuronal nitric oxide synthase and cFOS similar to those found in human schizophrenia *Neuroscience* 126(3): 591-598.
- Kirk RE (2012) *Experimental design: Procedures for the behavioral sciences* Thousand Oaks: Sage Publications
- Kishimoto J, Tsuchiya T, Cox H, Emson PC, Nakayama Y (1998) Age-related changes of calbindin_D18k, calretinin, and parvalbumin mRNAs in the hamster brain *Neurobiology of Aging* 19(1): 77-82.
- Kittelberger K, Hur EE, Sazegar S, Keshavan V, Kocsis B (2012) Comparison of the effects of acute and chronic administration of ketamine on hippocampal oscillations: relevance for the NMDA receptor hypofunction model of schizophrenia *Brain Structure and Function* 217(2): 395-409.
- Klausberger T (2009) GABAergic interneurons targeting dendrites of pyramidal cells in the CA1 area of the hippocampus *European Journal of Neuroscience* 30: 947-957.
- Knott GW, Quairiaux C, Genoud C, Welker E (2002) Formation of dendritic spines with GABAergic synapse induced by whisker stimulation in adult mice *Neuron* 34(2): 265-273.
- Kubicki M, McCarley R, Westin CF, Park HJ, Maier S, Kikinis R, Jolesz FA, Shenton ME (2007) A review of diffusion tensor imaging studies in schizophrenia *Journal of Psychiatric Research* 41: 15-30.
- Kukshal P, Thelma BK, Nimgaonkar VL, Deshpande SN (2012) Genetics of schizophrenia from a clinical perspective *International Review of Psychiatry* 24(5): 393-404.
- Kuruba R, Hattiangady B, Parihar VK, Shuai B, Shetty AK (2011) Differential susceptibility of interneurons expressing neuropeptide Y or parvalbumin in the aged hippocampus to acute seizure activity *PLoS ONE* 6(9): e24493.
- Leucht S, Corves C, Arbter D, Engel RR, Li C, Davis JM (2009) Second-generation versus first-generation antipsychotic drugs for schizophrenia: A meta-analysis *The Lancet* 373(9657): 31-41.

- Leuner B, Glasper ER, Gould E (2010) Sexual experience promotes adult neurogenesis in the hippocampus despite an initial elevation in stress hormones *PLoS ONE* 5(7): e11597.
- Lewis DA, Curley AA, Glausier JR, Volk DW (2012) Cortical parvalbumin interneurons and cognitive dysfunction in schizophrenia *Trends in Neuroscience* 35(1): 57-67.
- Lewis DA, Hashimoto T, Volk DW (2005) Cortical inhibitory neurons and schizophrenia *Nature Reviews Neuroscience* 6: 312-324.
- Lewis DA, Lieberman JA (2000) Catching up on schizophrenia: Natural history and neurobiology *Neuron* 28: 325-334.
- Lewis DA, Pierri JN, Volk DW, Melchitzky DS, Woo TUW (1999) Altered GABA neurotransmission and prefrontal cortical dysfunction in schizophrenia *Biological Psychiatry* 46(5): 616-626.
- Lewis DA, Sweet RA (2009) Schizophrenia from a neural circuitry perspective: Advancing toward rational pharmacological therapies *Journal of Clinical Investigation* 119(4): 706-716.
- Lewis DA, Volk DW, Hashimoto T (2004) Selective alterations in prefrontal cortical GABA neurotransmission in schizophrenia: A novel target for the treatment of working memory *Psychopharmacology* 174(1): 143-150.
- Lisman JE, Coyle JT, Green RW, Javitt DC, Benes FM, Heckers S, Grace AA (2008) Circuit-based framework for understanding neurotransmitted and risk gene interactions in schizophrenia *Trends in Neuroscience* 31(5): 234-242.
- Lolova I, Davidoff M (1992) Age-related morphological and morphometrical changes in parvalbumin- and calbindin-immunoreactive neurons in the rat hippocampal formation *Mechanisms of Ageing and Development* 66: 195-211.
- Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C (2004) Interneurons of the neocortical inhibitory system *Nature Reviews: Neuroscience* 5: 793-807.
- McBain CJ, Fisahn A (2001) Interneurons unbound *Nature Reviews: Neuroscience* 2: 11-23.
- Miettinen R, Sirviö J, Riekkinen P, Laakso MP, Riekkinen M, Riekkinen P (1993) Neocortical, hippocampal and septal parvalbumin- and somatostatin-

- containing neurons in young and aged rats: Correlation with passive avoidance and water maze performance *Neuroscience* 53(2): 367-378.
- Milner B, Squire LR, Kandel ER (1998) Cognitive neuroscience and the study of memory *Neuron* 20: 445-468.
- Morris BJ, Cochran SM, Pratt JA (2005) PCP: from pharmacology to modeling schizophrenia *Current Opinion in Pharmacology* 5(1): 101-106.
- Myer-Lindenberg AS, Olsen RK, Kohn PD, Brown T, Egan MF, Weinberger DR, Berman KF (2005) Regionally specific disturbance of dorsolateral prefrontal-hippocampal functional connectivity in schizophrenia *JAMA Psychiatry* 62(4): 379-386.
- Nitsch R, Soriano E, Frotscher M (1990) The parvalbumin-containing nonpyramidal neurons in the rat hippocampus *Anatomy and Embryology* 181(5): 413-425.
- Nomura T, Fukuda T, Aika Y, Heizmann CW, Emson PC, Kobayashi T, Kosaka T (1997a) Distribution of nonprincipal neurons in the rat hippocampus with special reference to their dorsoventral difference *Brain Research* 751(1): 64-80.
- Nomura T, Fukuda T, Aika Y, Heizmann CW, Emson PC, Kobayashi T, Kosaka T (1997b) Laminar distribution of non-principal neurons in the rat hippocampus, with special reference to their compositional difference among layers *Brain Research* 764: 197-204.
- Nowicka D, Soulsby S, Skangiel-Kramaska J, Glazewski S (2009) Parvalbumin-containing neurons, perineuronal nets and experience-dependent plasticity in murine barrel cortex *European Journal of Neuroscience* 30(11): 2053-2063.
- Ouda L, Druga R, Syka J (2008) Changes in parvalbumin immunoreactivity with aging in the central auditory system of the rat *Experimental Gerontology* 43(8): 782-789.
- Owen MJ (2012) Implications of genetic findings for understanding schizophrenia *Schizophrenia Bulletin* 38(5): 904-907.
- Patz S, Grabert J, Gorba T, Wirth MJ, Wahle P (2004) Parvalbumin expression in visual cortical interneurons depend on neuronal activity and TrkB ligands during an early period of postnatal development *Cerebral Cortex* 14(3): 342-351.

- Paxinos G, Watson C (1997) The rat brain in stereotaxic coordinates
NY:Academic Press.
- Pelayo-Terán JM, Suárez-Pinilla P, Chadi N, Crespo-Facorro B (2012) Gene-environment interactions underlying the effect of cannabis in first episode psychosis *Current Pharmaceutical Design* 18(32): 5024-5035.
- Perälä J, Suvisaari J, Saarni SI, Kuoppasalmi K, Isometsä E, Pirkola S, Partonen T, Tuulio-Henriksson A, Hintikka J, Kieseppä T, Härkänen T, Koskinen S, Lönngqvist J (2007) Lifetime prevalence of psychotic and bipolar I disorders in a general population *JAMA Psychiatry* 64(1): 19-28.
- Potier B, Jouvenceau A, Epelbaum J, Dutar P (2006) Age-related alterations of GABAergic input to CA1 pyramidal neurons and its control by nicotinic acetylcholine receptors in rat hippocampus *Neuroscience* 142: 187-201.
- Raghanti MA, Spocter MA, Butti C, Hof PR, Sherwood CC (2010) A comparative perspective on minicolumns and inhibitory GABAergic interneurons in the neocortex *Frontiers in Neuroanatomy* 4(3) 1-46.
- Romon T, Mengod G, Adell A (2011) Expression of parvalbumin and glutamic acid decarboxylase-67 after acute administration of MK-801: Implications for the NMDA hypofunction model of schizophrenia *Psychopharmacology* 217(2): 231-238.
- Schmitt A, Hasan A, Gruber O, Falkai P (2011) Schizophrenia as a disorder of disconnectivity *European Archives of Psychiatry and Clinical Neuroscience* 261: 150-154.
- Schobel SA, Kelly MA, Corcoran CM, Van Heertum K, Seckinger R, Goetz R, Harkavy-Friedman J, Malaspina D (2009) Anterior hippocampal and orbitofrontal cortical structural brain abnormalities in association with cognitive deficits in schizophrenia *Schizophrenia Research* 114: 110-118.
- Seto-Ohshima A, Emson PC, Berchtold MW, Heizmann CW (1989) Localization of parvalbumin mRNA in rat brain by in situ hybridization histochemistry *Experimental Brain Research* 75: 653-658.
- Shenton ME, Dickey CC, Frumin M, McCarley RW (2001) A review of MRI findings in schizophrenia *Schizophrenia Research* 49: 1-52.
- Shetty AK, Turner DA (1998) Hippocampal interneurons expressing glutamic acid decarboxylase and calcium-binding proteins decrease with aging in Fischer 344 rats *The Journal of Comparative Neurology* 394(2): 252-269.

- Sigmundsson T, Suckling J, Maier M, Williams SCR, Bullmore ET, Greenwood KE, Fukuda R, Ron MA, Toone BK (2001) Structural abnormalities in frontal, temporal, and limbic regions and interconnecting white matter tracts in schizophrenic patients with prominent negative symptoms *The American Journal of Psychiatry* 158(2): 234-243.
- Sik A, Penttonen M, Buzsaki G (1995) Hippocampal CA1 interneurons: An in vivo intracellular labeling study *The Journal of Neuroscience* 15(10): 6651-6665.
- Siucinska E, Kossut M (2006) Short-term sensory learning does not alter parvalbumin neurons in the barrel cortex of adult mice: a double-labeling study *Neuroscience* 138(2): 715-724.
- Sloviter RS (2004) Calcium-binding protein (calbindin-D28k) and parvalbumin immunocytochemistry: Localization in the rat hippocampus with specific reference to the selective vulnerability of hippocampal neurons to seizure activity *The Journal of Comparative Neurology* 280(2): 183-196.
- Stanley EM, Fadel JR, Mott DD (2012) Interneuron loss reduces dendritic inhibition and GABA release in hippocampus of aged rats *Neurobiology of Aging* 33(2): 431.e1-341.e13.
- Steullet P, Cabungcal JH, Kulak A, Kraftsik R, Chen Y, Dalton TP, Cuenod M, Do KQ (2010) Redox dysregulation affects the ventral but not dorsal hippocampus: Impairment of parvalbumin neurons, gamma oscillations, and related behaviors *Journal of Neuroscience* 30(7): 2547-2558.
- Stevens KE, Johnson RG, Rose GM (1997) Rats reared in social isolation show schizophrenia-like changes in auditory gating *Pharmacology Biochemistry and Behavior* 58(4): 1031-1036.
- Szilagy T, Orban-Kis K, Horvath E, Metz J, Pap Z, Pavai Z (2011) Morphological identification of neuron types in the rat hippocampus *Romanian Journal of Morphology & Embryology* 52: 15-20.
- Tamminga CA, Buchanan RW, Gold JM (1998) The role of negative symptoms and cognitive dysfunction in schizophrenia outcomes *International Clinical Psychopharmacology* 13(3): 21-26.
- Tan HY, Callicott JH, Weinberger DR (2007) Dysfunctional and compensatory prefrontal cortical systems, genes and the pathogenesis of schizophrenia *Cerebral Cortex* 17: 171-181.

- Uhlhaas PJ, Singer W (2010) Abnormal neural oscillations and synchrony in schizophrenia *Nature Reviews Neuroscience* 11: 100-113.
- Van der Borght K, Wallinga AE, Luiten PGM, Eggen BJL, Van der Zee EA (2005) Morris water maze learning in two rat strains increases the expression of the polysialylated form of the neural cell adhesion molecule in the dentate gyrus but has no effect on hippocampal neurogenesis *Behavioral Neuroscience* 119(4): 926-932.
- van Os J, Kenis G, Rutten BP (2010) The environment and schizophrenia *Nature; Perspectives* 468: 203-212.
- van Praag H, Kempermann G, Gage FH (2000) Neural consequences of environmental enrichment *Nature Reviews Neuroscience* 1: 191-198.
- Vela J, Gutierrez A, Vitorica J, Ruano D (2003) Rat hippocampal GABAergic molecular markers are differentially affected by ageing *Journal of Neurochemistry* 85: 368-377.
- Volk DW, Lewis DA (2002) Impaired prefrontal inhibition in schizophrenia: Relevance for cognitive dysfunction *Physiology and Behavior* 77:501-505.
- Volk DW, Matsubara T, Li S, Sengupta EJ, Georgiev D, Minabe Y, Sampson A, Hashimoto T, Lewis DA (2012) Deficits in transcriptional regulators of cortical parvalbumin neurons in schizophrenia *The American Journal of Psychiatry* 169(10): 1082-1091.
- Wang CZ, Yang SF, Xia Y, Johnson KM (2008) Postnatal phencyclidine administration selectively reduces adult cortical parvalbumin-containing interneurons *Neuropsychopharmacology* 33: 2442-2455.
- Wedzony K, Fijal K, Mackowiak M, Chocyk A (2008) Detrimental effect of postnatal blockade of N-methyl-D-aspartate receptors on sensorimotor gating is reversed by neuroleptic drugs *Pharmacological Reports* 60(6): 856-864.
- Williams SM, Goldman-Rakic PS, Leranth C (1992) The synaptology of parvalbumin-immunoreactive neurons in the primate prefrontal cortex *The Journal of Comparative Neurology* 320(3): 353-369.
- Woodruff AR, Anderson SA, Yuste R (2010) The enigmatic function of chandelier cells *Frontiers in Neuroscience* 4: 201.
- Woolley SC, Doupe AJ (2008) Social context-induced song variation affects female behavior and gene expression *PLoS Biology* 6(3): e62.

- Zhang Y, Behrens MM, Lisman JE (2008) Prolonged exposure to NMDAR antagonist suppresses inhibitory synaptic transmission in prefrontal cortex *Journal of Neurophysiology* 100: 959-965.
- Zhang ZJ, Reynolds GP (2002) A selective decrease in the relative density of parvalbumin-immunoreactive neurons in the hippocampus in schizophrenia *Schizophrenia Research* 55(1-2): 1-10.
- Zoladz PR, Park CR, Halonen JD, Salim S, Alzoubi KH, Srivareerat M, Fleshner M, Alkadhi KA, Diamond DM (2012) Differential expression of molecular markers of synaptic plasticity in the hippocampus, prefrontal cortex, and amygdala in response to spatial learning, predator exposure, and stress-induced amnesia *Hippocampus* 22(3): 577-589.
- Zuschratter W, Scheich H, Heizmann CW (1985) Ultrastructural localization of the calcium-binding protein parvalbumin in neurons of the song system of the zebra finch, *Poephila guttata* *Cell and Tissue Research* 241: 77-83.

Figures




Figure 1: *Location of representative sections relative to Bregma.*

PV-stained sections were pseudo-randomly selected for analysis of PV expression across the long axis of the hippocampus, somatosensory cortex, and retrosplenial cortex. Representative Nissl sections were used to help delineate areas and can be seen in this figure. Septal CA1, CA3, and DG samples, as well as retrosplenial and somatosensory (barrel field) cortex samples, were taken from sections within -2.45mm through -3.9mm relative to Bregma (A). Midseptotemporal CA1, CA3, and DG, in addition to temporal CA1 and CA3, samples were taken from sections within -5.0mm through -6.0mm relative to Bregma (B). Sampled sections were roughly 240 μ m apart from one another.




Figure 2: *Optimizing PV Antibody Concentration for Visualization of PV-containing Interneurons.*

In order to determine the optimal anti-parvalbumin antibody concentration, a subset of rats ($n = 2$) were used to compare three different concentrations: 1:1000 (A), 1:4000 (B), and 1:7000 (C). Based on qualitative observations, it was determined that an antibody concentration of 1:4000 was optimal for the purpose of visualizing PV expression for the current study. As evident in A, 1:1000 produced a high amount of background staining making it difficult to distinguish PV-expressing cell bodies. In contrast in C, a concentration of 1:7000 resulted in very light staining, making it difficult to distinguish whether some of the possible expression was, indeed, PV-expressing cells or simply staining artifacts.




Figure 3: *Anatomical locations of photomicrographs taken for analysis of PV expression in hippocampal regions of interest.*

Photomicrographs of septal hippocampal sub-regions were taken between areas were taken between -2.45mm through 3.9mm relative to Bregma **(A)**, while midseptotemporal and temporal sections were taken between -5mm through -5mm relative to Bregma **(B)**. Septal hippocampal photomicrographs of CA1 **(A.1)**, CA3 **(A.2)**, and DG **(A.3)** were taken at 20x magnification. Midseptotemporal CA1 **(B.1)**, CA3 **(B.2)**, and DG **(B.3)**, as well as temporal CA1 **(B.4)** and CA3 **(B.5)** were also taken at 20x magnification. Photomicrographs of all regions were taken within the boxed regions of interest as outlined above.




Figure 4: *Example of PV-expressing interneuron quantification using ImageJ color-based thresholding.*

Cell counting was automatized using ImageJ and region-specific macros in order to ensure unbiased, consistent quantification within each area of interest. An example of threshold-based quantification, in this case from hippocampal CA1, can be seen above. The first panel shows an original photomicrograph of CA1 with PV-expressing interneurons clearly evident, particularly within the pyramidal cell layer. The second panel shows the same photomicrograph, thresholded based on color density to isolate the most darkly stained cells. A macro was then run to count the number of PV-expressing cells based on previously set parameters and exclude any thresholded particles (e.g. non-cell artifacts) that do not meet criteria for inclusion (e.g. size, sphericity), and provided an output identifying the counted cells (as can be seen in the third panel).




Figure 5: *Septotemporal differences in PV expression as a function of hippocampal sub-region and age.*

6-month old rats had significant septotemporal variation in PV expression within hippocampal CA1 **(A)** and DG **(B)**, but not in CA3 **(C)**. The most striking difference in 6-month old rats were the decreases from septal to midseptotemporal, however, there was no difference between midseptotemporal and temporal CA1 in 6-month old rats **(A)**. There was no effect of septotemporal location in 1- and 24-month old rats in any hippocampal sub-region **(A-C)**.

* $p < .05$




Figure 6: *Distribution of HPC CA1 PV+ interneurons as a function of age and septotemporal location.*

Representative photomicrographs of hippocampal CA1 PV expression across age and septotemporal location. A clear age-related decline in PV-expression from 1- to 6-month old naïve rats in midseptotemporal and temporal CA1 can be seen even through qualitative comparison. 24-month old extensively trained rats have comparatively higher PV expression across all septotemporal levels of CA1.




Figure 7: *Distribution of HPC CA3 PV+ interneurons as a function of age and septotemporal location.*

Representative photomicrographs of hippocampal CA3 PV expression across age and septotemporal locations. There were no significant age-related declines in PV expression within any septotemporal level of CA3.




Figure 8: *Distribution of HPC DG PV+ interneurons as a function of age and septotemporal location.*

Representative photomicrographs of hippocampal DG PV expression across age and septotemporal location. From simple observation, it is clear that there are far more PV+ interneurons in extensively trained 24-month old DG. Furthermore, it is evident that there is a significant age-related decline in PV expression from 1- to 6-month old naïve rats.




Figure 9: Age-dependent decrease in parvalbumin expression within the hippocampal sub-regions CA1 and DG, but not CA3.

Within CA1 (**A**) there is a significant age-related decrease in PV expression from 1- to 6-month old rats in midseptotemporal and temporal locations. This age-related decline in PV expression is also seen in septal and midseptotemporal DG (**B**). There is no noncommittal decrease in PV expression in any septotemporal region in hippocampal CA3 (**C**). These data suggest that PV expression decreases within CA1 and DG as age increases. * $p < .05$




Figure 10: 24-month-old rats have significantly higher PV+ cells within retrosplenial and somatosensory cortices than 1- and 6-month-old rats.

Photomicrographs of the cortical regions of interest were taken from -2.45mm through -3.9mm relative to Bregma, with retrosplenial cortical photos taken at 20x magnification (**A.1**), and somatosensory (barrel field) cortical photos taken at 10x

magnification to ensure visualization of all cortical lamina (**A.2**). 24-month old extensively trained rats had significantly higher PV+ cells within the retrosplenial cortex (**B**) and somatosensory cortex (**C**) than both 1- and 6-month old rats. There was no statistically significant difference between PV expression within these cortices between 1- and 6-month old experimentally naïve rats. These findings suggest that extensive training increases PV+ expression within these cortical regions. $*p < .01$




Figure 11: *PV+ expression within the retrosplenial cortex across age groups.*

As shown statistically in Figure 10B, it is clear through visual observation that 24-month old extensively trained rats (**C**) have significantly more PV expression within the retrosplenial cortex than 1- (**A**) and 6- (**B**) month old naïve rats. Though there was no statistically different expression between 1 and 6 month old rats, the PV+ interneurons in 1 month old rats (**A**) appear visually larger than both 6 and 24 month old rats. It is unknown why, though this observation could be due to larger cell size in young rats, or perhaps an increase in PV expression.




Figure 12: *PV+ expression within the somatosensory (barrel field) cortex: distribution depends on age.*

Qualitative observations within the somatosensory cortex barrel fields revealed an age-dependent distribution of PV expression across cortical layers. The top panels (**A-C**) show full photomicrographs of the barrel field cortex for 1 (**A**), 6 (**B**), and 24 (**C**) month old rats. It is clear that 24 month old extensively trained rats have significantly more PV+ interneurons than both 1 and 6 month old naïve rats. Bottom panels (**D-F**) show higher magnification of top panels to illustrate laminar differences in PV distribution based on age of the animal. 1 month old rats have PV expression almost exclusively in the deep cortical layers (**D**), while 6 month (**E**) old rats (while having statistically similar overall expression to 1 month old rats) have PV expression in both deep and superficial cortical layers (but not in layers 3/4). In contrast, 24 month old rats show PV expression in all cortical layers containing cell bodies (**F**).






Figure 13: *Hippocampal CA1 has significantly more PV expression than CA3 and DG in septal and midseptotemporal regions across all age groups.*

Across all age groups (1-, 6-, and 24-months), regardless of past experience, CA1 PV expression was significantly higher in septal and midseptotemporal levels when compared to CA3 and DG **(A-C)**. Interestingly, in 24-month old extensively trained rats **(C)** there was no difference between CA1 and CA3 at the midseptotemporal level, though the DG had significantly lower PV expression than either CA1 or CA3. There was no difference between PV expression in temporal locations between CA1 and CA3 regions. * $p < .05$




Figure 14: 24-month old extensively trained rats have elevated PV expression across all hippocampal regions.

In CA1 and CA3, there were no septotemporal differences in PV expression in 24-month old rats; however, there was a significant decrease in PV expression between septal and midseptotemporal DG (**A**). Within CA1 (**B**), 24-month old

extensively trained rats had significantly higher PV expression than 6-month old naïve rats at all septotemporal levels. Furthermore, 24-month old rats had significantly higher PV expression than 1-month old rats in septal CA1, though there were no differences between the two groups in midseptotemporal or temporal CA1. Within the DG **(C)**, 1- and 24-month old rats had significantly higher PV expression than 6-month old rats in septal CA1. There were no differences between 1- and 24-month old rats at either level, and no difference between 1- and 6-month old rats in midseptotemporal DG, though a significant difference between 6- and 24-month old rats was seen in this region. 24-month old rats had significantly higher PV expression than both 1- and 6-month old rats in midseptotemporal and temporal CA3 regions **(D)**. No differences were seen between age-groups in septal CA3. * $p < .05$




Figure 15: *Proposed age-dependent decline in PV expression as a function of brain region.*

Here, we propose a theoretical framework for PV expression as a function of age and brain region. These are approximations based on the data collected in the present study, and reflect an age-dependent decline in PV expression that continues with age, as is seen in the 1-month to 6-month decrease in hippocampal, but not cortical, areas. We propose that over the course of the lifespan, hippocampal regions (especially CA1 and DG, based on our results) would be most impacted, and CA3 would be minimally impacted, while

neocortical regions would remain relatively stable over the lifespan. Circles indicate observations seen in 24-month old behaviorally trained rats in the current study, while dotted lines indicate proposed levels of PV expression we would expect to see in naïve aged rats.

Tables

Table 1. Average number of parvalbumin-positive interneurons across cortical and hippocampal regions in the rat brain as a function of age and septotemporal location.

| | | 1 month <i>M</i> ± <i>SEM</i> | <i>n</i> | 6 months <i>M</i> ± <i>SEM</i> | <i>n</i> | 24 months <i>M</i> ± <i>SEM</i> | <i>n</i> |
|---------------------------------|------------------|----------------------------------|----------|-----------------------------------|----------|------------------------------------|----------|
| <i>Cortex</i> | | | | | | | |
| Retrosplenial ^a | | 50.4±8.2 | 8 | 41.5±10.7 | 8 | 95.2±21.0 | 8 |
| Barrel Field ^b | | 73.5±18.1 | 8 | 105.4±22.0 | 8 | 250.9±31.8 | 8 |
| <i>Hippocampus</i> ^c | | | | | | | |
| CA1 | Septal | 15.6±2.2 | 8 | 12.1±1.7 | 8 | 19.6±2.1 | 8 |
| | Midseptotemporal | 12.0±1.3 | 8 | 5.9±1.3 | 8 | 15.6±2.3 | 8 |
| | Temporal | 13.5±2.4 | 7 | 4.3±0.9 | 8 | 17.1±1.4 | 8 |
| CA3 | Septal | 6.9±1.5 | 8 | 3.2±1.6 | 8 | 9.9±2.4 | 8 |
| | Midseptotemporal | 3.7±1.4 | 6 | 2.9±1.1 | 8 | 11.8±1.2 | 8 |
| | Temporal | 8.6±0.9 | 6 | 6.9±1.9 | 8 | 14.8±1.2 | 8 |
| DG | Septal | 5.0±0.5 | 8 | 0.6±0.3 | 8 | 6.8±1.5 | 8 |
| | Midseptotemporal | 2.4±1.0 | 6 | 0.3±0.2 | 8 | 3.6±1.2 | 8 |

^a analyzed region was 1600µm x 1000µm

^b analyzed region was 800µm²

^c analyzed region was 800µm x 500µm